Equilibrium dec 15 file 2

Monday, December 12, 2022 5:55 PM

Doubt Step 2

(11), (15), (17), (66), (13), (14), (69), 103, (51)

(11) $\Delta G^\circ = -RT \ln K_p$ $\boxed{K_p = P_{CO_2}}$

$-13.5\times10^3 = -2.303 \times R \times T \log(P_{CO_2})$ → 8.314 or $\frac{100}{12}$

(15) $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ $\frac{92\times20}{92 \times 100}$

t=0 1 mole 0

0.2

atm $1-0.2$ 0.4

0.8

% mass = % mole diss

Total moles at eq^m = 1.2

$PV = nRT$ (initially)

$1 \times V = 1 \times R \times 300$

$V = 300R$

$PV = nRT$

$P \times 300R = 1.2 \times R \times 600$

$\boxed{P = 2.4 \text{ atm}}$

Haber's Process

(17) $N_2 + 3H_2 \rightleftharpoons 2NH_3$; $\Delta H = -46.1$ kJ/mole

Catalyst ⇒ Scrap Fe or Fe oxide → Fe_2O_3, Fe_3O_4

Promoter (to ↑ efficiency of catalyst) ⇒ Molybdenum (Mo)

(66) $NH_4SH(s) \rightleftharpoons NH_3(g) + H_2S(g)$ → 3L, 600K

0.1 mole 0 0

$0.1 - 0.1\alpha$ 0.1α 0.1α

→ Total moles of Gas $= 0.2\alpha$

$= 0.2 \times \frac{30}{100}$

$= 0.06$

$K_c = [NH_3][H_2S]$

$= \left(\frac{0.1\alpha}{3}\right)^2$

$= \left(\frac{0.1\times0.3}{3}\right)^2 = 10^{-4} = \frac{K_p}{(RT)^{\Delta n_g}}$

$10^{-4} = \frac{K_p}{(0.0821 \times 600)^2}$

(69) $A(s) \rightleftharpoons B(g) + C(g)$; $K_{p_1} = x$ atm^2

P_1 P_1+P_2

$D(s) \rightleftharpoons C(g) + E(g)$; $K_{p_2} = y$ atm^2

P_2+P_1 P_2

→ $P_{Total} = P_B + P_C + P_E$

$= P_1 + P_1 + P_2 + P_2$

$= 2(P_1+P_2)$

$= 2(\sqrt{x+y})$

$x = K_{p_1} = P_1 \times (P_1+P_2)$

$y = K_{p_2} = P_2 \times (P_2+P_1)$

$K_{p_1} + K_{p_2} = x + y = (P_1+P_2)(P_1+P_2)$

$x+y = (P_1+P_2)^2$

$\sqrt{x+y} = (P_1+P_2)$

(*) Case of Incomplete neutralisation b/w $\boxed{W.A + S.B}$ / $\boxed{S.A + W.B}$

Case (a) equivalents of W.A > equivalents of S.B

Sol^n will have W.A + Salt in the container ; such Sol^n are called ACIDIC BUFFER

Case b: eq of W.B > eq of S.A

Sol^n contains W.B + Salt at the end of Rx^n; Such Sol^n is called BASIC Buffer.

Buffer Sol^n:- Sol^n in which negligible pH change is observed upon add^n of small amount of Acid or Base.

or

Sol^n which resist change in pH upon add^n of small add^n of acid or Base.

Acidic Buffer:- Mixture of W.A + its Salt from S.B or W.A + its Conjugate Base

eg (i) $CH_3COOH + CH_3COONa$ [$(CH_3COO)_2Ca$]

CH_3COO^- (C.B of CH_3COOH)

(ii) $HF | F^-$ (W.A C.B) or $HF + CaF_2$ or $HF + KF$ or $HF + NaF$ ~~$HF + NH_4F$~~

(iii) $(H_2CO_3 + NaHCO_3)$ or $(H_2CO_3 | HCO_3^-)$ (W.A C.B)

(iv) $(NaHCO_3 + Na_2CO_3)$ or $HCO_3^- | CO_3^{2-}$ → Composition of Blood buffer

(v) ~~$HCN + NH_4CN$~~

(vi) $HCN + NaCN$

$\underset{C-C\alpha}{CH_3COOH} \rightleftharpoons CH_3COO^-$ (Salt) $+ H^+$

$K_a = \frac{[CH_3COO^-]_{salt}[H^+]}{[CH_3COOH]}$

$[H^+] = K_a \times \frac{[W.A]}{[C.B]_{salt}}$ ⇒ $-\log[H^+] = -\log K_a + \log\frac{[C.B]}{[W.A]}$ (pH, pKa)

W.B | C.A

Q. 0.1M CH_3COOH ($pK_a = 4.7$) + 0.1M $(CH_3COO)_2Ca$, pH = ? → $pH = 4.7 + \log\left(\frac{0.2}{0.1}\right)$

pH Calculat^n of Acidic Buffer

eg $CH_3COOH + CH_3COONa$

$\alpha << 1$ ⇒ due to CIE, $C - C\alpha \approx C$

$CH_3COOH \rightleftharpoons CH_3COO^- + H^+$ (W.E)

$CH_3COONa \rightarrow CH_3COO^- + Na^+$ (S.E)

⇒ $K_a = \frac{[CH_3COO^-]_{Total}[H^+]}{[CH_3COOH]} = \frac{[CH_3COO^-]_{salt}[H^+]}{[CH_3COOH]}$

$[H^+] = K_a \frac{[CH_3COOH]}{[CH_3COO^-]_{salt}}$

$-\log[H^+] = -\log K_a - \log\frac{[CH_3COOH]}{[CH_3COO^-]_{salt}}$

$\boxed{pH = pK_a + \log\frac{[CH_3COO^-]_{salt}}{[CH_3COOH]}}$

(*) [weak acid] remains almost same due to CIE

Common ion

Due to CIE, α of CH_3COOH ↓es

$[CH_3COO^-]_{Total} = [CH_3COO^-]_{W.A} + [CH_3COO^-]_{salt} \approx [CH_3COO^-]_{salt}$

For Acidic Buffer

$pH = pK_a + \log\frac{[C.B \text{ of } W.A]}{[W.A]}$

or

$pH = pK_a + \log\frac{[\text{Anion of Salt}]}{[W.A]}$

Henderson-Hasselbalch Eq^n

(2) BASIC Buffer

→ Mix of Weak Base + its salt from S.Acid

or

W.B + its Conjugate Acid

eg i) $NH_4OH + NH_4Cl$ ($NH_3(aq)$) or $NH_3 | NH_4^+$ (aq)


(ii) $NH_4OH + (NH_4)_2SO_4$

(*) pOH of Basic Buffer $= pK_b + \log\frac{[\text{Cation of } W.B]}{[W.B]}$ or $pOH = pK_b + \log\frac{[C.Acid]}{[W.B]}$

$\boxed{pH = 14 - pOH}$

→ Henderson-Hasselbalch Eq^n.

Q. 100ml, 0.1M CH_3COOH ($pK_a = 4.774$) ; 100ml, 0.1M CH_3COONa



pH of Sol^n = ?

Sol^n:- Acidic Buffer

mmoles of $CH_3COOH = 100 \times 0.1$

$[CH_3COOH]_{new} = \frac{100\times0.1}{Total\ V} = \frac{100\times0.1}{200} = \frac{1}{20} = 0.05M$

$[CH_3COO^-]_{salt} = [CH_3COONa]_{new} = 0.05M$

$pH = pK_a + \log\frac{[CH_3COO^-]_{salt}}{[CH_3COOH]} = 4.774 + \log\frac{0.05}{0.05}$ (= 1)

$\boxed{pH = pK_a} = 4.774$

(3) Salt Buffer (It is a salt sol^n formed by Complete neutralizat^n of W.A + W.B)

eg CH_3COONH_4, NH_4CN, NH_4F, $(NH_4)_2CO_3$ etc.

(*) $\boxed{pH = 7 + \frac{1}{2}[pK_a - pK_b]}$

Q. find pH of Sol^n

1) 100ml, 0.1M CH_3COOH + 50ml, 0.1M NaOH ($pK_a = 4.774$)

2) $\frac{3}{4}$th of CH_3COOH is neutralised by NaOH

3) Half of CH_3COOH is neutralised by NaOH → pH = pKa

H.W 4) 100ml, 0.1M CH_3COOH + 200ml, 0.1M NaOH (10mmole, 20mmole)

(1) $CH_3COOH(aq) + NaOH(aq) \rightarrow CH_3COONa(aq) + H_2O(l)$

t=0 10mmole 5mmole 0 –

-5 -5 $+5$

5 0 5

$[CH_3COOH] = \frac{5}{150}M$

$[CH_3COONa] = \frac{5}{150}M = [CH_3COO^-]$ → Acidic Buffer

$[W.A] = [C.B]$ ⇒ $\boxed{pH = pK_a}$

(2) $\frac{1}{4}$th CH_3COOH remaining, $\frac{3}{4}$th CH_3COONa formed

$pH = pK_a + \log\frac{(3/4)}{(1/4)}$

$pH = 4.77 + \log 3$

$= 4.774 + 0.47$

(4) $CH_3COOH + NaOH \rightarrow CH_3COONa + H_2O(l)$

10 20 0 –

-10 -10 $+10$ –

0 10 10

$[NaOH]_{Strong} = \frac{10}{300}M$

$[CH_3COONa] = \frac{10}{300}M$

→ $CH_3COO^- + H_2O \rightleftharpoons CH_3COOH + OH^-$

$Na^+ + H_2O \rightarrow \times$

$[OH^-]_{Total} = [OH^-]_{NaOH} + [OH^-]$ (Anionic hydrolysis of CH_3COO^-) → negligible

$[OH^-]_{Total} \approx [OH^-]_{Strong\ Base} = \frac{10}{300} = \frac{1}{30}M$

$pOH = -\log\left(\frac{1}{30}\right)$

$\boxed{pH = 14 - pOH}$

Buffer Action

(I) let Few drops of Strong Acid (HCl) added to Acidic Buffer ($CH_3COOH + CH_3COONa$)

$\log\frac{[C.B]}{[W.A]} = \log\frac{[CH_3COO^-]}{[CH_3COOH]}$

$CH_3COO^- + H^+ \rightleftharpoons CH_3COOH$ (C.B, added Acid, W.A)

$[C.B]$↓, $[W.A]$↑, $\frac{[C.B]}{[W.A]}$ ↓es

$pH = pK_a + \log\frac{[C.B]}{[W.A]}$ → ↓es

pH ↓es negligibly

(II) Small amount OH^- is added to $(HCN + CN^-)$ Acidic Buffer

pH ↑es negligibly $[HCN]$↓ $[CN^-]$↑

(III) Small amount of H^+ added $(NH_4OH + (NH_4)_2SO_4)$ BASIC Buffer


pH↓, pOH↑, $[NH_4OH]$ ↓es, $[NH_4^+]$ ↑es

(H.W) Example 13, 14, 15, 16 (Pg - 134, 135) → V.V. important

---

Solubility of Salt and Ksp

max amount of Salt that can be dissolved to achieve Saturat^n at a given T in given amount of solvent



NaCl (Saturated Sol^n) → Brine

Very less soluble Salt in H_2O Called Sparingly soluble salt

$AgCl(solid) \rightleftharpoons Ag^+(aq) + Cl^-(aq)$

a − s mol/L s s

Solubility of Salt ⇓ Molarity

$K_{eq} = K_{sp} = [Ag^+][Cl^-]$

$K_{sp} = s^2$ Solubility product $\boxed{s = \sqrt{K_{sp}}}$

(#) $PbCl_2(s) \rightleftharpoons Pb^{2+}(aq) + 2Cl^-(aq)$

s 2s

s = solubility of $PbCl_2$

$K_{sp} = [Pb^{2+}][Cl^-]^2$

$= s \times (2s)^2$

$= 4s^3$

$s = \left(\frac{K_{sp}}{4}\right)^{1/3}$

(#) $A_xB_t \rightleftharpoons xA^{y+} + tB^{x-}$

a − s xs ts

Ag_2S_3 → $2Ag^{3+} + 3S^{2-}$

$K_{sp} = [A^{t+}]^x [B^{x-}]^t$

$= (xs)^x (ts)^t$

$\boxed{K_{sp} = x^x t^t s^{(x+t)}}$